

Rheumatism

No 34

#2

A  
Thesis

On Rheumatism.



By Joseph K Swift of Penna

read March 28 1816

In proper  
Papers  
collected, as the

any new i.

it has ab

Physicians

its best h.

a selection

would allow

Rheum

Class of Py

His divi

and Ebro.

con, with

sep. for w

ten in 14

shall

Acute

common

In proposing myself a candidate for a Medical Degree, I  
selected, as the subject of my Thesis. That I should advance  
any new ideas upon it, can scarcely be expected, after  
it has already been treated of by so many eminent  
Physicians; I therefore assert no pretensions to original-  
ity, but have merely endeavoured to make as judicious  
a selection from the writings of others as my abilities  
would allow. —

Rheumatism has been arranged by Dr Cullen in his  
Class of Pyrexia, and Order of Phlegmasia. —

It is divided into two distinct species, viz the Acute  
and Chronic. It is the acute Rheumatism only, which  
can, with propriety, be said to belong to the Phlegma-  
sia, for we discover little, or no inflammatory ac-  
tion in the Chronic stage. —

I shall commence with the consideration of the  
Acute Rheumatism. It is one of the most  
common diseases in all climates where a great

*[Faint, illegible handwriting on the left page, likely bleed-through from the reverse side.]*

capitula  
from the  
only fo  
the great  
and Vern  
an sim  
y ar.  
From the  
tion an  
with ear  
tinct, an  
the mos  
Rheum  
and is m  
Rheum  
Joint.  
Rheum  
as the h



vicissitude of weather occurs, but in those of an uniform temperature, whether warm or cold, it is more rarely found: In this country, we generally observe the greatest number of cases during the autumnal and Vernal months, but we see it prevailing, under similar circumstances, at every season of the year.

From the resemblance of their symptoms, Rheumatism and Gout have frequently been confounded with each other; they are however specifically distinct, and it may not be improper to designate the most material points wherein they differ. Rheumatism is less confined to the feet than Gout, and is more liable to shift its place. —

Rheumatic pains are not so shooting as those of Gout. —

Rheumatism principally affects the larger joints, as the knee, hip &c. whereas Gout attacks the

*[Faint, illegible handwriting on the left page, likely bleed-through from the reverse side.]*

Rheumat  
Just i  
Rheum  
stomach  
Rheum  
turn at  
The Bre  
most su  
Rheum  
bet Gour  
In Rhe  
matter  
are move  
Other p  
my mig  
amiated  
between  
Rheum

✓ The smaller ones. — Rheumatism is not hereditary  
Gout is —

✓ Rheumatism is not preceded by affections of the  
stomach, or other Viscera, as Gout is. —

Rheumatism is not, like Gout, disposed to re-  
turn at stated intervals. —

✓ The British Physicians say that women are  
most subject to Rheumatism; men, to Gout. —

Rheumatism may occur at any period of life,  
but Gout is usually confined to adults. —

In Rheumatism, the disposition of calcareous  
matter in the joints so frequently observed in Gout,  
are never discovered. —

Other particulars in which the two diseases vary  
might be mentioned, but those already enu-  
merated, will enable us to form a diagnosis  
between them. —

Rheumatism is found to prevail most in marsh

sily, an  
tency  
Rugine  
The Re  
ly occur  
part of  
Bamph  
the feet  
Cullen  
miasm  
is in m  
the Br  
Mecho  
rise to  
ers, po  
In son  
to the  
be Bar

soils, and Dr Barton has remarked, that it is extremely common in some calcareous counties in Virginia. -

The Remote causes are various. It is most frequently occasioned by the application of cold to one part of the body, whilst the others are kept warm. Dampness, from wearing wet garments, or getting the feet wet, oftentimes produces it. -

Cullen says Rheumatism proceeds from Marsh Miasmata, and of this there can be no doubt, as in many instances we perceive it putting on the Intermittent form.

Mechanical matters of various kinds may give rise to it, and to this its frequency among painters, potters and gilders may be ascribed. -

In some particular cases it has been attributed to the fumes of Arsenic. -

Dr Barton related the case of a young gentleman

*[Faint, illegible handwriting on the left page, likely bleed-through from the reverse side.]*

in a  
prop  
gent in  
ted wo  
the dis  
mation  
It may  
gry of  
Mlay  
The O  
He din  
clino of  
and A  
The Y  
an gen  
someti  
from on  
the kn

in whom it originated from the sudden suppression of the gonorrhoeal discharge by astringent injections; every medicine which was exhibited was ineffectual, and afforded no relief until the discharge was restored, when the Rheumatism disappeared.

It may arise from the suppression of Hemorrhaging of any kind, and from the healing up of Ulcers, and some cutaneous affections, as the Stole.

The Predisposing Causes are Cold— a variable climate— the age of puberty and the decline of life— excessive evacuations— Spring and Autumn.— great muscular irritation.

The Symptoms are— pains of the joints— these are generally confined to the joints alone, but sometimes affect the muscles in their passage from one joint to another. The larger joints, as the knee, hip and shoulder, are most liable

to be the  
felt in  
toms of  
fused,  
which  
sympt  
When  
monly  
derable  
time.  
The pr  
joint, a  
turn a  
transl  
The ex  
ring the  
Permit  
the pa



to be the seats of the disease. The pains are mostly felt in particular parts, before any febrile symptoms appear, and when they are extensively diffused, the disease is ushered in by a cold fit, which is immediately succeeded by the other symptoms. —

When there is little or no fever the pain is commonly confined to one joint; but if it be considerable, several may be affected at the same time. —

The pains do not commonly remain long in one joint, but shift to others, and occasionally return again to parts from which they had been translated. —

The exacerbations of the fever take place during the night, observing an Intermittent or Remittent form, and it is during this period that the pains generally shift. —

*[Faint, illegible handwriting on the left page, likely bleed-through from the reverse side.]*

After  
society  
The p  
but no  
Of sw  
commo  
very c  
The p  
one is  
of the  
gives  
The b  
Glam  
Thoug  
Pleas  
supp  
mita  
These

After the joints have been affected for some time, they swell, and become red and painful to the touch. The pains are mostly mitigated by this swelling, but not always entirely removed. —

If sweating come on, it commonly does so in the commencement of the disease, but it is seldom very copious, nor does it often prove critical. —

The pulse is full, hard and frequent. The urine is high-coloured, and in the early stage of the disease, without sediment, but as it progresses, it deposits the luteous sediment. —

The blood drawn from a vein exhibits the inflammatory crust. —

Though Rheumatism differs from most of the Plethorisms in scarcely ever terminating in suppuration, yet effusions of a serous fluid are sometimes made into the sheaths of the tendons. These tumours should not be opened but left

for the  
them p  
cult to  
With  
continu  
in some  
it may  
Abund  
The ease  
of the t  
ven for  
It is re  
limp a  
Mr. Dr  
of a hor  
On dig  
disease,  
The tra

for the absorbent vessels to remove, as incisions into them produce ulcers, which are extremely difficult to heal.

With these symptoms Rheumatism generally continues for fourteen or fifteen days; though in some cases, even under the best management it may continue for several weeks.

Rheumatism of the Intermittent kind, though the easiest to cure with proper care, is frequently of the longest duration, sometimes remaining even for years. -

It is seldom a dangerous disease, but it sometimes attacks the heart, when it is mostly fatal. Mr. Brindley had nine patients thus affected, of whom, seven died. -

On dissection of persons who have died of this disease, effusions are at times found within the cranium, or topical affections of some of the

visera  
cases  
of a w  
of the n  
effusio  
Poc  
imate  
to be  
other  
crease  
is app

In the  
for in  
secute  
Vena  
of the  
one of

viscera are perceived; in some of the above mentioned cases of Mr Dundas, the heart was enlarged, and of a whitish colour. — In the joints, thickening of the membranes, adhesions, and gelatinous effusions are discovered. —

Respecting the different theories of the Proximate cause, I shall say nothing. According to Dr Barton it is the same as that of any other inflammation, depending upon an increased afflux of blood to a part, whilst it is exposed to the action of cold. —

### Cure of Acute Rheumatism

In the curative treatment, the usual remedies for inflammation must be vigorously prosecuted.

Venesection. From the inflammatory nature of the disease, it must be obvious, that this is one of the most important remedy. The blood

must  
get, and  
public  
demand  
time, be  
extent  
incur  
one to  
The co  
thoug  
iliary  
It is of  
and an  
The or  
large,  
in as  
Topica  
oftent



must be drawn largely during the first stages, and the bleeding, repeated as often as the pulse and the state of the symptoms may demand: we must, however, at the same time, be cautious not to carry it to such an extent as to induce debility, and thereby incur the danger of bringing on the Chronic stage.

The cure should not be trusted to it alone, though it is one of the most powerful auxiliary means. —

It is of least service in Intermittent cases, and in that species proceeding from miasmata. The orifice made into the vein should be large, that the blood may be evacuated in as short a space of time as possible. Topical bleeding by cupping or leeches is oftentimes beneficial, so here the pain of

The joy  
ridings  
a cur,  
Nery  
the inf  
Eathu  
irritate  
However  
letting  
and epl  
with c  
of the p  
ing to s  
In some  
marked  
at the  
ced, an  
toms of

The joints is very acute, and where there is much redness and swelling. But it seldom effects a cure, and is very liable to shift the pains. -

Purgings is of great importance in lessening the inflammatory state of the system. Those Cathartics which produce the least general irritation, are to be preferred. They do not, however, reduce the fever so speedily as Blood-letting; and where the disease is very severe and extensive, their exhibition is attended with considerable inconvenience, on account of the pain the patient experiences in going to stool. -

In some particular cases there exists a remarkable excitement of the arterial system at the same time, the patient is much reduced, and labours under many of the symptoms of Chronic Rheumatism. For the remo-

val of  
inade  
long a  
Dia  
ficiency  
they  
mini  
temp  
Durin  
laid a  
be att  
whic  
most  
ly re  
from  
munt  
stant  
four

val of this excitement, Venisection has been found inadequate, but it has entirely ceased under long continued purging with Sulphur.

Diaphoretics are medicines of considerable efficacy in Rheumatism. In very acute cases they are improper, and should not be administered until the inflammatory symptoms have been reduced by depletion.

During their exhibition, the general rules laid down for conducting sweating should be attended to. Those of a mild nature, and which occasion the least irritation, are the most proper. The Dover's Powder is most strongly recommended, but to obtain any benefit from it, 'tis necessary to continue its employment for some time, and to keep up a constant perspiration with it for at least twenty four hours, and as some direct, for a much longer

period  
in no  
ter ble  
stoma  
with  
times  
Nig  
from  
rial  
that  
minis  
to Ba  
adver  
Nite  
cy and  
the  
decrea  
mause

period. Nitre is said to have been serviceable in many cases. It should be given shortly after bleeding, in doses as large as the patient's stomach will bear. It is likewise combined with Tart. Emet. Jams Powder, and sometimes with Digitalis.

Digitalis has not been much employed, and from the vast influence it has over the arterial system, we might suppose, a priori, that it would be of great importance in diminishing inflammatory action.

Dr Barton has given it, he says, with much advantage, either alone or in union with Nitre, effecting a reduction of the frequency and fullness of the pulse, as well as of the symptoms generally. Its virtues are not decreased by its occasioning some degree of nausea. Yet, notwithstanding, as its opera-

tion co  
and a  
in the  
it sho  
Em  
be adm  
found  
Intern  
M  
cite  
of acc  
ended  
La sy  
in inc  
The O  
ion is  
tion.  
by other



tion cannot be, at all times uniformly obtained and as the promptest remedies are required in the commencement of the disease, I think it should never be substituted for U.S. purging.

Emetics, if the stomach be in a foul state, may be administered with much benefit. They are found most useful in Rheumatism of an Intermittent type.

Mercurials are exhibited with a view to excite Ptyalism and purging. In many cases of acute Rheumatism, a salivation is succeeded by the happiest effects, and in those of a syphilitic nature, Mercury is always an indispensable article. —

The Preparations of Arsenic, a diversity of opinion is entertained of their virtues in Rheumatism. By some, their efficacy is maintained; by others, it is denied. Dr. Barton says that in

many  
if such  
nature  
early  
effecti  
seen i  
einsy  
guage  
imur  
demon  
ious, c  
eino  
Whene  
coince  
Pauv  
stage.  
tend  
Butter

many cases, Arsenic acts as a specific, and that it succeeds best in those of the most acute nature. He also remarks that it is particularly useful in Intermittent Rheumatism affecting the joints and Thorax, where he has seen it perform cures, when all other medicines had failed. This, however is the language of enthusiasm, and numerous experiments made with Arsenic in this City have demonstrated fully, that it is, at best, a precarious, and too frequently, a dangerous medicine in this disease. —

Whenever it is of any service, its good effects are evinced in a short time. —

Rauvian Bark is improper in the inflammatory stage. It is especially adapted to the Intermittent form, and should be given during the Intermission. —

*[Faint, illegible handwriting on the left page]*

Blasted  
when  
mently  
csmu  
tion of  
tion to  
confine  
ly bene  
Cold  
may be  
The Me  
dusie  
jed in  
purpo  
I shan  
tion.  
In de  
acute a

Blisters and Rubefacients, in very acute cases when the pains are severe and not permanently fixed, may, perhaps, be injurious, in as much as they tend to produce a translation of them to other parts: but after depletion has been promised, and the pains are confined to particular joints, they are highly beneficial. —

Cold Applications to the inflamed parts may be made with safety and advantage. The Russian Physicians are in the habit of using snow or pounded ice — Cloths dipped in cold water or Sac. Sat. answer every purpose. — — — —

I shall now treat of the Chronic Rheumatism. —

In drawing a line of distinction between the acute and Chronic stages, I shall make use

of the  
and still  
some  
of hyp  
specially  
disease  
the nat  
when tr  
ing; as  
redness  
they can  
while a  
out on  
old on  
ally, the  
old an  
the cas  
in Rh

of the language of Dr Eullen: When the pains are still ready to shift their place, when at the same time they are attended with some degree of pyrexia, and with some swelling, and especially, with some redness of the joints, the disease is to be considered as partaking of the nature of the acute Rheumatism. But when there is no degree of pyrexia remaining; when the pained joints are without redness, when they are cold and stiff; when they cannot be easily made to sweat, or when, while a free and warm sweat is brought out on the rest of the body, it is clammy and cold on the pained joints; and when, especially, the pains of the joints are increased by cold and relieved by heat applied to them, the case is to be considered as purely Chronic Rheumatism." The differences between

*[Faint, illegible handwriting on the left page of an open manuscript.]*

the  
sai  
pa  
Ab  
que  
tion  
erac  
stag  
an  
wh  
to su  
usid  
to the  
twine  
disca  
and  
The a  
nt the



the two forms of the disease are here very accurately delineated, with the exception of one particular. Cullen says, that in Chronic Rheumatism there is no pyrexia; but we frequently perceive a slight degree of febrile action remaining for some time after the general characteristic symptoms of the Chronic stage have supervened. Dr Barton proposed an amendment of this part of his definition which, to me, appears extremely judicious, viz. to substitute in the place of *Nulla*, the term used by Dr Cullen, that of *Subnulla pyrexia*. -

Dr Rush also had an intermediate grade between the two forms, as his division of the disease into *Rheumatismus*, *Rheumaticula*, and *Rheumatalgia*, implied. -

The Acute form is generally sooner relieved by art than the Chronic. -

...the ...  
...but ...  
...those ...  
...of me ...  
...and ...  
The ...  
The ...  
of the ...  
by is ...  
The ...  
by ...  
our ...  
turned ...  
The ...  
prepa ...  
it is, ...  
is goi ...  
sing ...

Chronic Rheumatism may be seated in any joint, but it is more particularly disposed to infest those which are surrounded with a number of muscles and integuments, as the Hip joint and the Vertebrae.

#### The Cure of Chronic Rheumatism.

The general indication is to restore the energy of the vital principle in the diseased parts, by internal and external remedies.

The different Preparations of Guaiacum may be exhibited with the greatest advantage. They are sometimes proper before the disease has attained the Chronic form.

The Volatile Tincture is preferable to the other Preparations. The best method of employing it is, to give a large dose of it when the patient is going to bed.  $\mathfrak{zss}$  or even  $\mathfrak{zj}$  - at the same time using some diluent drink, as Wine &c. to

*[Faint, illegible handwriting on the left page, likely bleed-through from the reverse side.]*

apist  
in sub  
The O  
their p  
The V  
Butter  
county  
infla  
until  
to op  
rum  
ned co  
give, i  
nj or 3p  
An po  
The Sa  
st su  
of Rhe

assist its operation. The Guaiac: may be given in substance in doses of 10 or 15 grs. —

The Balsams are nearly allied to Guaiac: in their properties. —

The Volatile Alkali has been administered by the British Physicians in the acute stage. In this country, however, the disease is always of a too inflammatory nature to allow its exhibition until Venisection &c. have been premised. In its operation it seems nearly allied to the Guaiacum, but is inferior to it. It may be combined with the Guaiac: or when this is offensive, it may be given alone, to the amount of ʒij or ʒj per day. —

In particular cases of Chronic Rheumatism the Savin has been exhibited with the greatest success. The discrimination of those forms of Rheumatism, which require its use, demands

*[Faint, illegible handwriting on the left page, likely bleed-through from the reverse side.]*

consider  
the be  
there is  
insure  
attended  
has been  
It's pie  
It invig  
minist  
It prom  
in Syph  
nation  
It sho  
invea  
fully  
Profy  
win ex  
privat

considerable attention. Where the surface of the body is cold, with clammy sweats, when there is a debility of the joints, frequently amounting to an almost entire loss of motion, attended with acute topical pain, the Sarsaparilla has been given with the most salutary effect. Its primary operation is that of a stimulant, it invigorates the force of the circulation, diminishes the pain, and excites perspiration. It promises, likewise to be a valuable medicine in Syphilitic Rheumatism, and in Rheumatism, complicated with Gout. —

It should be given in doses of 10 or 15 grs. and increased, more or less, till its effects are fully developed. —

Professor Eschschman has employed the Sarsaparilla extensively in his public, as well as private practice, and does not hesitate to

prom  
in C  
Can  
Have  
Unbr  
spoon  
The  
to, b  
unly  
long  
Lup  
phys  
Of an  
ing  
whole  
alwa  
When  
the



pronounce it superior to all other medicines in Chronic Rheumatism. —

Cantbarides, in Substance and in Tinctures, have been advantageously employed.

Unbruised Mustard-seeds, in doses of a Table spoonful, have been found serviceable. —

The Preparations of Iron have been resorted to, but little benefit is derived from them unless their exhibition is continued for a length of time. —

Turpentine has been recommended by some Physicians. —

If any febrile action should remain, purging will be beneficial, and through the whole course of the disease, costiveness should always be obviated. —

When the symptoms continue refractory, and the various medicines resorted have proved

insufficient  
a slight  
naged  
If the  
severe,  
The O  
Carto  
effect  
m. al  
wise,  
trap  
In so  
confi  
Cicut  
prepa  
a. clat  
imm  
Where

25  
insufficient, advantage has been derived from  
a slight salivation, but it requires to be ma-  
naged with the greatest caution. —

If the symptoms be urgent, and the pains  
severe, Opiates may be given to procure rest.

The *Phytolacca Bicandra*, according to Dr  
Barton, has been administered with good  
effect in those cases where *Guaiacum* is  
ineffectual. In Syphilitic Rheumatism, like-  
wise, in combination with Calomel, its vir-  
tues have been considerable. —

In some Syphilitic cases of a Chronic nature  
benefit has resulted from the employment of  
*Picuta* in combination with the mercurial  
preparations — it relieves pain, and produces  
a determination to the surface. —

Among the External remedies for Chronic  
Rheumatism, Blisters, and other stimulating

apple  
Blist  
or the  
by son  
Cabin  
Acco  
more  
ed join  
to ple  
disca  
over t  
The p  
and  
mus  
Cold  
bun  
The  
direct

applications are entitled to our attention. The Blisters should be either frequently repeated or the irritation caused by them continued by some stimulant ointment, as the Ung. Sabin. or of Cantharides. —

According to Dr Thomas it will be found more useful, in certain cases where the affected joints are surrounded with large muscles, to place the blister at some distance from the diseased parts, than to lay it immediately over them. —

The patient should be clothed in flannel, and flannel shirts substituted for linen or muslin ones. —

Cold water, by affusion, or the cold bath has been highly recommended by some physicians. The Warm Bath, or the Vapor of warm water, directed to the part, has frequently proved use-

July 18<sup>th</sup>  
 City of  
 New York  
 Dear  
 Friend  
 I have  
 the pleasure  
 to inform  
 you that  
 the  
 same  
 has been  
 received  
 and  
 forwarded  
 to  
 the  
 proper  
 authorities  
 for  
 their  
 consideration  
 and  
 approval  
 I am  
 very  
 respectfully  
 Yours  
 Wm. L. Garrison

ful. It diminishes the pain, and relaxes the rigidity of the muscular fibres. —

Friiction of the parts with flannel or the flesh brush should always be enjoined. —

Exercise of the part, or of the whole body, in any moderate manner, is of service. —

The Diet should be cordial and nutritious, and the patient may drink Wine whey, or Barley water, in which Elix crystal of Tartar have been dissolved.

In cases where a great degree of muscular contraction occurs, and the joints are in danger of becoming permanently flexed, to obviate the inconvenience resulting from a bent limb, and to prevent deformity, Dr Balfour, of Edinburgh, has advised the application of a splint to the diseased part, so as to keep it in a state of extension. —

With this, Gentlemen, I conclude my subject, and am fully sensible that many apologies are necessary.

up for the  
 treated.  
 minded  
 want of  
 led those  
 useful a  
 if in t  
 myself  
 and a  
 my m  
 edoe, r



ing for the imperfect manner in which it has been  
treated. There are many remedies, highly recom-  
mended, which I have been compelled to omit for  
want of room. I have, however, endeavoured to se-  
lect those, which appeared to me particularly  
useful in the management of the disease; but  
if, in this selection, I have erred, or have shown  
myself deficient in perspicuity of language  
and arrangement, I trust that you will attribute  
my mistakes to inexperience and want of knowl-  
edge, rather than to obstinacy and prejudices.